

STELLATE GANGLION BLOCKS

Stellate ganglion blocks are utilized to anesthetize the sympathetic nerve fibers in the neck. Patients who are candidates for this procedure suffer from sympathetically mediated pain, generally known as complex regional pain or reflex sympathetic dystrophy, in the upper extremities and sometimes the head and neck. To ensure proper placement of the medication, the procedure is performed under fluoroscopic imaging, a low dose type x-ray.

These injections usually contain local anesthetic medications. If effective, they can be repeated several times to ensure maximum pain relief.

Generally, these procedures are performed in the surgery center. The patient will be asked to change into a hospital gown. An IV will be started for medications and safety. A brief pre-procedure history and physical examination are performed. All patient medications taken at home will be reviewed to ensure there are no contraindications to the procedure and that the patient has complied with pre-procedure instructions. The patient will be required to have a complete list of all medication normally taken at home. The patient is then transported to the procedure room and positioned on an x-ray table. The skin will then be cleaned with a sterile soap and draped in a sterile fashion. The skin over the injection target site is then anesthetized with a local anesthetic. Fluoroscopy is then used to guide a needle into the proper location. Contrast dye is administered (unless the patient has allergies to iodine) to confirm placement and the medication is slowly injected. Once the procedure is completed, the patient returns to the recovery area for approximately 30 minutes to rest.

The entire process for check-in, preparation, recovery and check-out will take 1-2 hours. The procedure itself generally lasts for only 15 minutes, but the rest of the time is necessary to ensure a safe, efficacious procedure. During the injection, pain is sometimes increased temporarily.

Every medical procedure, no matter how minor, contains some risks. Anytime, a needle is placed into the body there is a risk of tissue injury, infection and bleeding. If this occurs near the spine, it can result in nerve damage. Although extremely rare, nerve damage and death have occurred following spine injections. Injections near the spine also risk puncturing the dura and can result in headaches, which can easily be treated. More common risks include a temporary increase in pain, local tenderness after the injection, and allergic reactions to the medications.